

Homological stability of symplectic groups

Ismael Sierra

Nathalie Wahl

University of Toronto

Copenhagen University

Introduction: what is homological stability?

Homological Stability I: motivation

$G =$ (discrete) group.

$BG =$ classifying space of principal G -bundles.

$H^*(BG) =$ characteristic classes of such bundles.

Homological Stability I: motivation

$G =$ (discrete) group.

$BG =$ classifying space of principal G -bundles.

$H^*(BG) =$ characteristic classes of such bundles.

Goal: to compute $H^*(BG)$, or compute $H_*(BG)$.

Algebraic interpretation: $H_*(BG) = \text{Tor}_*^{\mathbb{Z}[G]}(\mathbb{Z}, \mathbb{Z})$ and

$H^*(BG) = \text{Ext}_{\mathbb{Z}[G]}^*(\mathbb{Z}, \mathbb{Z})$.

Homological Stability I: motivation

$G =$ (discrete) group.

$BG =$ classifying space of principal G -bundles.

$H^*(BG) =$ characteristic classes of such bundles.

Goal: to compute $H^*(BG)$, or compute $H_*(BG)$.

Algebraic interpretation: $H_*(BG) = \text{Tor}_*^{\mathbb{Z}[G]}(\mathbb{Z}, \mathbb{Z})$ and
 $H^*(BG) = \text{Ext}_{\mathbb{Z}[G]}^*(\mathbb{Z}, \mathbb{Z})$.

Problem: this is very hard!

Homological Stability II: a (partial) solution

Key idea (Quillen): Many groups of interest arise in families:

$$G_0 \hookrightarrow G_1 \hookrightarrow G_2 \hookrightarrow \dots .$$

Thus, get a family of classifying spaces

$$BG_0 \rightarrow BG_1 \rightarrow BG_2 \rightarrow \dots .$$

Then, we can ask two questions.

Homological Stability II: a (partial) solution

Key idea (Quillen): Many groups of interest arise in families:

$$G_0 \hookrightarrow G_1 \hookrightarrow G_2 \hookrightarrow \dots .$$

Thus, get a family of classifying spaces

$$BG_0 \rightarrow BG_1 \rightarrow BG_2 \rightarrow \dots .$$

Then, we can ask two questions.

- (i) Do we have *homological stability*? i.e. are the maps $H_d(BG_n) \rightarrow H_d(BG_{n+1})$ isomorphisms for $d \ll n$?
- (ii) Can we compute the *stable homology*? i.e. can we compute $\operatorname{colim}_n H_d(BG_n)$?

Homological Stability II: a (partial) solution

Key idea (Quillen): Many groups of interest arise in families:

$$G_0 \hookrightarrow G_1 \hookrightarrow G_2 \hookrightarrow \dots$$

Thus, get a family of classifying spaces

$$BG_0 \rightarrow BG_1 \rightarrow BG_2 \rightarrow \dots$$

Then, we can ask two questions.

- (i) Do we have *homological stability*? i.e. are the maps $H_d(BG_n) \rightarrow H_d(BG_{n+1})$ isomorphisms for $d \ll n$?
- (ii) Can we compute the *stable homology*? i.e. can we compute $\operatorname{colim}_n H_d(BG_n)$?

If answer to both is yes then we get partial computations!... and improving stability range becomes relevant!

Stable homology: examples

Fact: In many examples of interest, we can access the stable homology $H_d(BG_\infty) := \operatorname{colim}_n H_d(BG_n)$ using *group completion theorem*.

Stable homology: examples

Fact: In many examples of interest, we can access the stable homology $H_d(BG_\infty) := \operatorname{colim}_n H_d(BG_n)$ using *group completion theorem*.

Examples:

- (i) $G_n = GL_n(\mathbb{F}_q)$: stable homology completely described by Quillen '72.
- (ii) $G_n = GL_n(R)$: stable homology related to $K(R) =$ algebraic K -theory of R .
Example: $R =$ number field and \mathbb{Q} coefficients then known by Borel '74.
- (iii) $G_n = MCG(\Sigma_{n,1})$: stable homology known!
 $H^*(BG_\infty, \mathbb{Q}) = \mathbb{Q}[\kappa_1, \kappa_2, \dots]$. (Madsen–Weiss '07)
- (iv) $G_n = Sp_{2n}(\mathbb{Z})$: $H^*(BSp_\infty(\mathbb{Z}), \mathbb{Q}) = \mathbb{Q}[x_2, x_6, x_{10}, \dots]$ by Borel '74.
Integral computations possible by 9 authors (Calmès–Dotto–Harpaz–Hebestreit–Land–Moi–Nardin–Nikolaus–Steimle).

The “Classical” argument I: set-up

Question: how does one prove homological stability? for which families?

“Classical” argument (Quillen, Randal-Williams–Wahl,...) gives a framework for examples.

The “Classical” argument I: set-up

Question: how does one prove homological stability? for which families?

“Classical” argument (Quillen, Randal-Williams–Wahl,...) gives a framework for examples.

Set-up:

1. $(\mathcal{C}, \oplus, \mathbf{o}) =$ (braided) monoidal category.
2. $X \in \mathcal{C} =$ *stabilizing object*.
3. $A \in \mathcal{C} =$ choice of object, usually take $A = \mathbf{o}$.
4. $G_n = \text{Aut}_{\mathcal{C}}(A \oplus X^{\oplus n})$.
5. $G_n \hookrightarrow G_{n+1}$ given by $- \oplus \text{id}_X$.

The Classical argument II: examples

- (i) $F = \text{field}$, take $C = \text{Vect}_F^{f \cdot d}$, $\oplus = \text{direct sum}$, $X = F$, $A = 0$. Then $G_n = GL_n(F)$.

The Classical argument II: examples

(i) $F = \text{field}$, take $C = \text{Vect}_F^{f.d}$, $\oplus = \text{direct sum}$, $X = F$, $A = 0$. Then $G_n = GL_n(F)$.

(ii) $C = \begin{cases} \text{objects} = & \Sigma_{n,1} \\ \text{morphisms} = & \text{diffeomorphisms rel boundary/isotopy} \end{cases}$,



$\oplus = \text{boundary connected sum}$,



$A = \emptyset$, $X = \Sigma_{1,1}$. Then $G_n = MCG(\Sigma_{n,1})$.

The Classical argument II: examples

(i) $F = \text{field}$, take $C = \text{Vect}_F^{f.d}$, $\oplus = \text{direct sum}$, $X = F$, $A = 0$. Then $G_n = GL_n(F)$.

(ii) $C = \begin{cases} \text{objects} = & \Sigma_{n,1} \\ \text{morphisms} = & \text{diffeomorphisms rel boundary/isotopy} \end{cases}$,



$\oplus = \text{boundary connected sum}$,



$A = \emptyset$, $X = \Sigma_{1,1}$. Then $G_n = MCG(\Sigma_{n,1})$.

(iii) $C = \text{category of skew-symmetric bilinear forms over } \mathbb{Z}$, $\oplus = \text{orthogonal direct sum}$, $A = 0$, $X = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ hyperbolic form.
Then $G_n = Sp_{2n}(\mathbb{Z})$.

The Classical argument III: the method

There are canonical **destabilization complexes** $W_n(A, X)$ associated to the above data, dimension = $n - 1$.

Informally: p -simplices \longleftrightarrow ways of destabilizing $X^{\oplus p+1}$ from $A \oplus X^{\oplus n}$.

The Classical argument III: the method

There are canonical **destabilization complexes** $W_n(A, X)$ associated to the above data, dimension = $n - 1$.

Informally: p -simplices \longleftrightarrow ways of destabilizing $X^{\oplus p+1}$ from $A \oplus X^{\oplus n}$.

Theorem (Quillen, Randal-Williams–Wahl): If the connectivity of $W_n(A, X)$ grows with n then the family $\{G_n\}_n$ has homological stability.

The Classical argument III: the method

There are canonical **destabilization complexes** $W_n(A, X)$ associated to the above data, dimension = $n - 1$.

Informally: p -simplices \longleftrightarrow ways of destabilizing $X^{\oplus p+1}$ from $A \oplus X^{\oplus n}$.

Theorem (Quillen, Randal-Williams–Wahl): If the connectivity of $W_n(A, X)$ grows with n then the family $\{G_n\}_n$ has homological stability.

Precise: If $W_n(A, X)$ is $\frac{n-c}{k}$ connected, $c \in \mathbb{Z}, \in \mathbb{Z}_{>0}$ then $H_d(BG_n) \rightarrow H_d(BG_{n+1})$ is an iso if $d \leq \frac{n-c+2}{\max\{2, k\}}$.

The Classical argument III: the method

There are canonical **destabilization complexes** $W_n(A, X)$ associated to the above data, dimension = $n - 1$.

Informally: p -simplices \longleftrightarrow ways of destabilizing $X^{\oplus p+1}$ from $A \oplus X^{\oplus n}$.

Theorem (Quillen, Randal-Williams–Wahl): If the connectivity of $W_n(A, X)$ grows with n then the family $\{G_n\}_n$ has homological stability.

Precise: If $W_n(A, X)$ is $\frac{n-c}{k}$ connected, $c \in \mathbb{Z}, \in \mathbb{Z}_{>0}$ then $H_d(BG_n) \rightarrow H_d(BG_{n+1})$ is an iso if $d \leq \frac{n-c+2}{\max\{2, k\}}$.

Limitation of method: we can never get something better than $\frac{n-\text{const}}{2}$.

The Classical method: applications

- (i) $G_n = GL_n(R)$: get result $\frac{n-c}{2}$, $c = \text{constant}$ depending on ring. (Quillen, Maazen, Van der Kallen, ...).
- (ii) $G_n = MCG(\Sigma_{n,1})$ (Harer, Ivanov, Boldsen, Randal-Williams, Galatius–Kupers–Randal-Williams, Harr-Vistrup–Wahl). Best bound $\lesssim \frac{2n}{3}$.
- (iii) $G_n = Sp_{2n}(R)$ (Charney, Miraii–Van der Kallen,...) get range $d \lesssim \frac{n}{2}$, constant depends on ring.

The Classical method: applications

- (i) $G_n = GL_n(R)$: get result $\frac{n-c}{2}$, $c =$ constant depending on ring. (Quillen, Maazen, Van der Kallen, ...).
- (ii) $G_n = MCG(\Sigma_{n,1})$ (Harer, Ivanov, Boldsen, Randal-Williams, Galatius–Kupers–Randal-Williams, Harr-Vistrup–Wahl). Best bound $\lesssim \frac{2n}{3}$.
- (iii) $G_n = Sp_{2n}(R)$ (Charney, Miraii–Van der Kallen,...) get range $d \lesssim \frac{n}{2}$, constant depends on ring.

Surprise: bound for MCG is $2n/3 \gg n/2$... so hard to get using “classical method” alone...

The Classical method: applications

- (i) $G_n = GL_n(R)$: get result $\frac{n-c}{2}$, $c =$ constant depending on ring. (Quillen, Maazen, Van der Kallen, ...).
- (ii) $G_n = MCG(\Sigma_{n,1})$ (Harer, Ivanov, Boldsen, Randal-Williams, Galatius–Kupers–Randal-Williams, Harr–Vistrup–Wahl). Best bound $\lesssim \frac{2n}{3}$.
- (iii) $G_n = Sp_{2n}(R)$ (Charney, Miraii–Van der Kallen,...) get range $d \lesssim \frac{n}{2}$, constant depends on ring.

Surprise: bound for MCG is $2n/3 \gg n/2$... so hard to get using “classical method” alone...

Result of Harr–Vistrup–Wahl manages to do so!

Results

Theorem (S., Whal)

Let R be a ring with finite unitary stable rank (usr). Let $c = 0$ if R is a PID and $c = 2usr(R) + 2$ otherwise. Then

$$H_d(BSp_{2g}(R)) \rightarrow H_d(BSp_{2(g+1)}(R))$$

is an iso for $d \leq \frac{2g-c-2}{3}$.

Theorem (S., Whal)

Let R be a ring with finite unitary stable rank (usr). Let $c = 0$ if R is a PID and $c = 2usr(R) + 2$ otherwise. Then

$$H_d(BSp_{2g}(R)) \rightarrow H_d(BSp_{2(g+1)}(R))$$

is an iso for $d \leq \frac{2g-c-2}{3}$.

This improves previously $1/2$ slope bound to $2/3$.

Theorem (S., Whal)

Let R be a ring with finite unitary stable rank (usr). Let $c = 0$ if R is a PID and $c = 2usr(R) + 2$ otherwise. Then

$$H_d(BSp_{2g}(R)) \rightarrow H_d(BSp_{2(g+1)}(R))$$

is an iso for $d \leq \frac{2g-c-2}{3}$.

This improves previously $1/2$ slope bound to $2/3$.

As we will see, $2/3$ slope is related to MCG $2/3$ slope...

Result II

Define the “odd” symplectic groups $Sp_{2g+1}(R) := \text{Stab}_{Sp_{2g+2}(R)}(e_1)$,
where standard basis is $e_1, f_1, \dots, e_g, f_g$.

Have $Sp_0(R) \subset Sp_1(R) \subset Sp_2(R) \subset Sp_3(R) \subset \dots$.

Result II

Define the “odd” symplectic groups $Sp_{2g+1}(R) := \text{Stab}_{Sp_{2g+2}(R)}(e_1)$, where standard basis is $e_1, f_1, \dots, e_g, f_g$.

Have $Sp_0(R) \subset Sp_1(R) \subset Sp_2(R) \subset Sp_3(R) \subset \dots$.

Theorem (S., Whal)

Let R be a ring with finite unitary stable rank (usr). Let $c = 0$ if R is a PID and $c = 2usr(R) + 2$ otherwise. Then

$$H_d(BSp_n(R)) \rightarrow H_d(BSp_{n+1}(R))$$

is an iso for $d \leq \frac{n-c-3}{3}$.

Result II

Define the “odd” symplectic groups $Sp_{2g+1}(R) := \text{Stab}_{Sp_{2g+2}(R)}(e_1)$, where standard basis is $e_1, f_1, \dots, e_g, f_g$.

Have $Sp_0(R) \subset Sp_1(R) \subset Sp_2(R) \subset Sp_3(R) \subset \dots$.

Theorem (S., Whal)

Let R be a ring with finite unitary stable rank (usr). Let $c = 0$ if R is a PID and $c = 2usr(R) + 2$ otherwise. Then

$$H_d(BSp_n(R)) \rightarrow H_d(BSp_{n+1}(R))$$

is an iso for $d \leq \frac{n-c-3}{3}$.

New slope = $1/3$.

The proof

The geometric idea I

Key idea (Harr–Vistrup–Wahl) $\Sigma_{n+1,1}$ can be obtained by attaching **two** handles to $\Sigma_{n,1}$.



Thus, we can define new family by attaching **one** handle at a time:
 $\Sigma_{0,1}, \Sigma_{0,2}, \Sigma_{1,1}, \Sigma_{1,2}, \Sigma_{2,1}, \Sigma_{2,2}, \dots$

Old stability problem has twice the speed \Rightarrow Suffices stability of slope $1/3$ for new family!

The geometric idea I

Key idea (Harr–Vistrup–Wahl) $\Sigma_{n+1,1}$ can be obtained by attaching **two** handles to $\Sigma_{n,1}$.



Thus, we can define new family by attaching **one** handle at a time:
 $\Sigma_{0,1}, \Sigma_{0,2}, \Sigma_{1,1}, \Sigma_{1,2}, \Sigma_{2,1}, \Sigma_{2,2}, \dots$

Old stability problem has twice the speed \Rightarrow Suffices stability of slope $1/3$ for new family!

Issues:


1. One has to be careful attaching handles!
2. How to create a nice categorical set-up?



The geometric idea II

Take \mathcal{C} = category of *bidecorated surfaces*.


Objects = (Σ, I_0, I_1) .



The geometric idea II

Take $C =$ category of *bidecorated surfaces*.

Objects = (Σ, I_0, I_1) .




Morphisms = isotopy classes of diffeomorphisms preserving the intervals.

The geometric idea II

Take \mathcal{C} = category of *bidecorated surfaces*.

Objects = (Σ, I_0, I_1) .




Morphisms = isotopy classes of diffeomorphisms preserving the intervals.

Monoidal structure: glue along “half of the intervals”, and use the remaining halves to define the new intervals in the glued surface.

The geometric idea II

Take \mathcal{C} = category of *bidecorated surfaces*.

Objects = (Σ, I_0, I_1) .



Morphisms = isotopy classes of diffeomorphisms preserving the intervals.

Monoidal structure: glue along “half of the intervals”, and use the remaining halves to define the new intervals in the glued surface.


$A = \emptyset$.

$X = \text{disc}$.

The geometric idea II

Take \mathcal{C} = category of *bidecorated surfaces*.

Objects = (Σ, I_0, I_1) .



Morphisms = isotopy classes of diffeomorphisms preserving the intervals.

Monoidal structure: glue along “half of the intervals”, and use the remaining halves to define the new intervals in the glued surface.

$A = \emptyset$.

$X = \text{disc}$.

This solves both problems!

The geometric idea III

$W_n(\emptyset, X) = \text{complex of disordered arcs.}$

Vertices: non-separating arcs from b_0 to b_1 (up to isotopy).

p - simplex: collection $\{a_0, \dots, a_p\}$ of non-separating pairwise disjoint arcs such that orders at b_0, b_1 agree.

The geometric idea III

$W_n(\emptyset, X) =$ complex of disordered arcs.

Vertices: non-separating arcs from b_0 to b_1 (up to isotopy).

p -simplex: collection $\{a_0, \dots, a_p\}$ of non-separating pairwise disjoint arcs such that orders at b_0, b_1 agree.

Theorem (Harr-Vistrup-Wahl): W_n is $\frac{n-5}{3}$ -connected.

The geometric idea III

$W_n(\emptyset, X) = \text{complex of disordered arcs.}$

Vertices: non-separating arcs from b_0 to b_1 (up to isotopy).

p -simplex: collection $\{a_0, \dots, a_p\}$ of non-separating pairwise disjoint arcs such that orders at b_0, b_1 agree.

Theorem (Harr-Vistrup-Wahl): W_n is $\frac{n-5}{3}$ -connected.

This implies stability result of slope $2/3$ for $MCG(\Sigma_{g,1})$.

Formed spaces with boundary I

Key insight: Action on homology (with R coefficients) gives a map $MCG(\Sigma_{g,1}) \rightarrow Sp_{2g}(R)$. Want to find $Sp_{2g+1}(R)$ with maps $MCG(\Sigma_{g,2}) \rightarrow Sp_{2g+1}(R)$. Then study new family.

Formed spaces with boundary I

Key insight: Action on homology (with R coefficients) gives a map $MCG(\Sigma_{g,1}) \rightarrow Sp_{2g}(R)$. Want to find $Sp_{2g+1}(R)$ with maps $MCG(\Sigma_{g,2}) \rightarrow Sp_{2g+1}(R)$. Then study new family.

Better wish: find algebraic analogue of bidecorated surfaces and a functor from bidecorated surfaces to it.

Formed spaces with boundary I

Key insight: Action on homology (with R coefficients) gives a map $MCG(\Sigma_{g,1}) \rightarrow Sp_{2g}(R)$. Want to find $Sp_{2g+1}(R)$ with maps $MCG(\Sigma_{g,2}) \rightarrow Sp_{2g+1}(R)$. Then study new family.

Better wish: find algebraic analogue of bidecorated surfaces and a functor from bidecorated surfaces to it.

Solution: Category F_∂ of formed spaces with boundary.

- Objects = (M, λ, ∂) , $M =$ f.g. free R -module, $\lambda =$ skew-symmetric bilinear form, $\partial : M \rightarrow R$.
- Morphisms = module maps preserving λ, ∂ .

- Monoidal structure $\#$:

$$(M_1, \lambda_1, \partial_1) \# (M_2, \lambda_2, \partial_2) = \left(M_1 \oplus M_2, \begin{pmatrix} \lambda_1 & \partial_1^T \partial_2 \\ -\partial_2^T \partial_1 & \lambda_2 \end{pmatrix}, \partial_1 + \partial_2 \right).$$

- $X = (R, 0, \text{id})$.

Formed spaces with boundary II

Geometric interpretation: Functor from bidecorated surfaces to F_∂ defined by $(\Sigma, I_0, I_1) \mapsto (H_1(\Sigma \cup_{I_0 \sqcup I_1} H), \lambda, \partial)$ where $\partial : H_1(\Sigma \cup_{I_0 \sqcup I_1} H) \cong H_1(\Sigma, I_0 \sqcup I_1) \rightarrow \tilde{H}_0(I_0 \sqcup I_1) \cong R$ is boundary map.




Figure 1: Two bidecorated surfaces (Σ, I_0, I_1) and their associated surface $\Sigma^+ = \Sigma \cup H$

Formed spaces with boundary II

Geometric interpretation: Functor from bidecorated surfaces to F_∂ defined by $(\Sigma, I_0, I_1) \mapsto (H_1(\Sigma \cup_{I_0 \sqcup I_1} H), \lambda, \partial)$ where $\partial : H_1(\Sigma \cup_{I_0 \sqcup I_1} H) \cong H_1(\Sigma, I_0 \sqcup I_1) \rightarrow \tilde{H}_0(I_0 \sqcup I_1) \cong R$ is boundary map.




Figure 1: Two bidecorated surfaces (Σ, I_0, I_1) and their associated surface $\Sigma^+ = \Sigma \cup H$

Note: bidecorated disc $\mapsto X = (R, o, \text{id})$.

Formed spaces with boundary II

Geometric interpretation: Functor from bidecorated surfaces to F_∂ defined by $(\Sigma, I_0, I_1) \mapsto (H_1(\Sigma \cup_{I_0 \sqcup I_1} H), \lambda, \partial)$ where $\partial : H_1(\Sigma \cup_{I_0 \sqcup I_1} H) \cong H_1(\Sigma, I_0 \sqcup I_1) \rightarrow \tilde{H}_0(I_0 \sqcup I_1) \cong R$ is boundary map.




Figure 1: Two bidecorated surfaces (Σ, I_0, I_1) and their associated surface $\Sigma^+ = \Sigma \cup H$

Note: bidecorated disc $\mapsto X = (R, o, \text{id})$.

Functor is monoidal! This gives geometric meaning to $\#$.

Algebraic meaning uses $(M_1 \oplus M_2)^\vee \cong M_1^\vee \oplus M_2^\vee$ and $\Lambda^2(M_1 \oplus M_2)^\vee \cong \Lambda^2 M_1^\vee \oplus M_1^\vee \otimes M_2^\vee \oplus \Lambda^2 M_2^\vee$.

Even and odd symplectic groups

Now, natural to consider $G_n = \text{Aut}(X^{\#n})$.

Fact: $n = 2g + 1$ one gets $G_n \cong \text{Sp}_{2g}(R)$.

Even and odd symplectic groups

Now, natural to consider $G_n = \text{Aut}(X^{\#n})$.

Fact: $n = 2g + 1$ one gets $G_n \cong \text{Sp}_{2g}(R)$.

When $n = 2g$ one gets $G_n \cong \text{Stab}_{\text{Sp}_{2g}(R)}(e_1) = \text{Sp}_{2g-1}(R)$.

Thus, $G_n = \text{Sp}_{n-1}(R)$... even and odd are exchanged for us!

Even and odd symplectic groups

Now, natural to consider $G_n = \text{Aut}(X^{\#n})$.

Fact: $n = 2g + 1$ one gets $G_n \cong \text{Sp}_{2g}(R)$.

When $n = 2g$ one gets $G_n \cong \text{Stab}_{\text{Sp}_{2g}(R)}(e_1) = \text{Sp}_{2g-1}(R)$.

Thus, $G_n = \text{Sp}_{n-1}(R)$... even and odd are exchanged for us!

Fun fact: There is a braiding in full subcategory generated by X so get $B_n \rightarrow \text{Aut}(X^{\#n}) = \text{Sp}_{n-1}(R)$... this is (reduced) Bureau representation!

Destabilization complex and arcs I

Geometrically $W_n =$ disordered arc complex.

Question: what happens algebraically?

Destabilization complex and arcs I

Geometrically $W_n =$ disordered arc complex.

Question: what happens algebraically?

Arc:= $a \in M$ such that $\partial a = 1$.

Geometric meaning: geometric arc gives algebraic arc!

Destabilization complex and arcs I

Geometrically $W_n =$ unordered arc complex.

Question: what happens algebraically?

Arc: $a \in M$ such that $\partial a = 1$.

Geometric meaning: geometric arc gives algebraic arc!

Non-separating: $a \in M$ arc is *non-separating* if $\{\lambda(a, -), \partial\}$ unimodular in M^\vee .

Meaning: condition says there is $b \in M$ such that $\partial b = 1$ and $\lambda(a, b) = 1$: “ b connects the two sides of a ”.

Destabilization complex and arcs I

Geometrically $W_n =$ unordered arc complex.

Question: what happens algebraically?

Arc: $a \in M$ such that $\partial a = 1$.

Geometric meaning: geometric arc gives algebraic arc!

Non-separating: $a \in M$ arc is *non-separating* if $\{\lambda(a, -), \partial\}$ unimodular in M^\vee .

Meaning: condition says there is $b \in M$ such that $\partial b = 1$ and $\lambda(a, b) = 1$: “ b connects the two sides of a ”.

Jointly non-separating: $\{\lambda(a_0, -), \dots, \lambda(a_p, -), \partial\}$ unimodular in M^\vee .

Destabilization complex and arcs II

What about ordering condition?

We say that a_0, \dots, a_p are *disordered* if we can pick an ordering of them such that $\lambda(a_i, a_j) = 1$ for $i < j$.




Figure 2: Two disordered arcs crossing once inside the handle

Destabilization complex and arcs II

What about ordering condition?

We say that a_0, \dots, a_p are *disordered* if we can pick an ordering of them such that $\lambda(a_i, a_j) = 1$ for $i < j$.




Figure 2: Two disordered arcs crossing once inside the handle

Thus, get *algebraic disordered arc complex* $D(M, \lambda, \partial)$.

Vertices= non-separating arcs.

p -simplex: jointly non-separating and disordered.

Destabilization complex and arcs II

What about ordering condition?

We say that a_0, \dots, a_p are *disordered* if we can pick an ordering of them such that $\lambda(a_i, a_j) = 1$ for $i < j$.




Figure 2: Two disordered arcs crossing once inside the handle

Thus, get *algebraic disordered arc complex* $D(M, \lambda, \partial)$.

Vertices= non-separating arcs.

p -simplex: jointly non-separating and disordered.

Key fact: W_n and $D(X^{\#n})$ agree (on a skeleton).

Destabilization complex and arcs II

What about ordering condition?

We say that a_0, \dots, a_p are *disordered* if we can pick an ordering of them such that $\lambda(a_i, a_j) = 1$ for $i < j$.




Figure 2: Two disordered arcs crossing once inside the handle

Thus, get *algebraic disordered arc complex* $D(M, \lambda, \partial)$.

Vertices= non-separating arcs.

p -simplex: jointly non-separating and disordered.

Key fact: W_n and $D(X^{\#n})$ agree (on a skeleton).

Theorem (S.-Wahl) $D(M, \lambda, \partial)$ is $\frac{n-c}{3}$ -connected, where $c = 5$ if R is a PID and $c = 2usr(R) + 6$ in general.

Theorem (S.-Wahl) $D(M, \lambda, \partial)$ is $\frac{n-c}{3}$ -connected, where $c = 5$ if R is a PID and $c = 2usr(R) + 6$ in general.

Using the above one proves stability theorem!

Proof based on a “bad simplex argument”.

Finishing the proof

Proof based on a “bad simplex argument”.

Start with complex of all non-separating arcs, and try to deform maps (this is analogue to geometric proof!).

Finishing the proof

Proof based on a “bad simplex argument”.

Start with complex of all non-separating arcs, and try to deform maps (this is analogue to geometric proof!).

Key: Complex of non-separating arcs related to unimodular vectors complexes... that has connectivity of slope 1 in fact!

Key algebraic ingredient: understanding how X -genus decreases when we cut algebraic arcs... problem is that the X -genus (algebraic version of “number of handles”) generally drops by 2 and not by 1, that causes slope $1/3$ and not $1/2$.

Further possible works and applications

1. In the geometric arc complex, slope $1/3$ stability is optimal. What about in the algebraic arc complex? (nothing known...)
2. Use this to get a classical proof $2/3$ -slope stability for diffeomorphism groups of some high-dimensional manifolds.
3. What about quadratic symplectic groups? issue is the non-separating arc complex... all other steps work analogously and stability of slope $> 1/4$ has new geometric implications!
4. Can one use similar methods to improve the slope $1/4$ connectivity in the paper “Uniform twisted homological stability” by Miller–Patz–Petersen–Randal-Williams? Maybe go to slope $1/3$? (Ideal conjecture says it is $1/2$ and connects to number theory)